



ELSEVIER

Contents lists available at ScienceDirect

Neuropsychologia

journal homepage: www.elsevier.com/locate/neuropsychologia

Partial maintenance of auditory-based cognitive training benefits in older adults



Samira Anderson^{a,b,1}, Travis White-Schwoch^{a,b}, Hee Jae Choi^{a,b}, Nina Kraus^{a,b,c,d,e,*}

^a Auditory Neuroscience Laboratory, Northwestern University, Evanston, IL 60208, USA²

^b Communication Sciences, Northwestern University, Evanston, IL 60208, USA

^c Institute for Neuroscience Northwestern University Interdepartmental Neuroscience Program, Chicago, IL 60611, USA

^d Department of Neurobiology and Physiology, Northwestern University, Evanston, IL 60208, USA

^e Department of Otolaryngology, Northwestern University, Chicago, IL 60611, USA

ARTICLE INFO

Article history:

Received 13 January 2014

Received in revised form

27 July 2014

Accepted 30 July 2014

Available online 8 August 2014

Keywords:

Aging

Training

Auditory plasticity

Neural timing

Speech-in-noise recognition

Memory

Temporal processing

Frequency-following response (FFR)

ABSTRACT

The potential for short-term training to improve cognitive and sensory functions in older adults has captured the public's interest. Initial results have been promising. For example, eight weeks of auditory-based cognitive training decreases peak latencies and peak variability in neural responses to speech presented in a background of noise and instills gains in speed of processing, speech-in-noise recognition, and short-term memory in older adults. But while previous studies have demonstrated short-term plasticity in older adults, we must consider the long-term maintenance of training gains. To evaluate training maintenance, we invited participants from an earlier training study to return for follow-up testing six months after the completion of training. We found that improvements in response peak timing to speech in noise and speed of processing were maintained, but the participants did not maintain speech-in-noise recognition or memory gains. Future studies should consider factors that are important for training maintenance, including the nature of the training, compliance with the training schedule, and the need for booster sessions after the completion of primary training.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

In response to growing interest in the potential for training to remediate perceptual and cognitive deficits (Pichora-Fuller & Levitt, 2012), software developers have marketed a wide variety of training programs to older adults with concerns about declines in cognitive function and the ability to hear in noisy environments. The adaptability of computer technology allows for fine manipulation of training stimuli, such as consonant–vowel (CV) transition times in a speech signal, allowing users to progress from easy to challenging perceptual tasks. In a previous study, we used an auditory-based cognitive training program (described in Smith et al., 2009) that combined adaptively expanding CV transition

times and memory demands to determine if training improves neural and behavioral speech-in-noise processing and cognitive functions of memory and speed of processing in older adults (ages 55–70) (Anderson, White-Schwoch, Parbery-Clark, & Kraus, 2013a). We evaluated neural speech-in-noise processing with the frequency following response (FFR) to a CV syllable presented in quiet and noise and found that training decreased FFR peak latencies and peak variability, two putative measures of subcortical neural synchrony, and improved untrained measures of sentence recognition in noise, short-term memory, and speed of processing. We used the FFR because it reflects temporal processing deficits in older adults (Anderson, Parbery-Clark, White-Schwoch, & Kraus, 2012; Clinard & Tremblay, 2013; Vander Werff & Burns, 2011) and because it can be modulated by training in young adults (Carcagno & Plack, 2011; Song, Skoe, Banai, & Kraus, 2012; Song, Skoe, Wong, & Kraus, 2008). The FFR reflects neural transcription of stimulus properties and is highly modulated by cognitive influences (Kraus & Chandrasekaran, 2010; Krishnan, Gandour, & Bidelman, 2010). In the current study, we investigated whether or not these effects of training on FFR processing and perceptual and cognitive function were maintained in older adults six months after cessation of training. In the following paragraphs,

* Corresponding author at: Communication Sciences, Neurobiology & Physiology, Otolaryngology Auditory Neuroscience Laboratory, Northwestern University, 2240 Campus Drive, Evanston, IL 60208, USA. Tel.: +1 847 491 3181; fax: +1 847 491 2523.

E-mail address: nkraus@northwestern.edu (N. Kraus).

¹ Current address: Department of Hearing and Speech Sciences, University of Maryland, 0100 LeFrak Hall, College Park, MD 20742, USA.

² <http://www.brainvolts.northwestern.edu>.

we outline training considerations for older adults and the evidence for generalization to untrained tasks and persistence of training benefits.

Effective training should target declines in sensory and cognitive processing and other domains that are known to decline with age. Reductions in speed of processing may underlie broad changes in cognitive function, including memory (Salthouse, 1996). In the auditory domain, these declines may contribute to the older adult's difficulties when trying to understand spoken communication, especially in challenging listening situations. Precise temporal resolution in the auditory system is required for the discrimination of speech stimuli (Phillips, Gordon-Salant, Fitzgibbons, & Yeni-Komshian, 2000), both in individuals with hearing loss and with audiometrically-normal hearing. Examples of age-related deficits in temporal resolution are seen in delayed midbrain and cortical peak latencies (Anderson et al., 2012; Clinard & Tremblay, 2013; Lister, Maxfield, Pitt, & Gonzalez, 2011; Tremblay, Piskosz, & Souza, 2003), delayed recovery in single neurons in inferior colliculus (Walton, Frisina, & O'Neill, 1998), loss of temporal coding by fusiform cells in dorsal cochlear nucleus (Wang et al., 2009), lower response amplitudes to amplitude-modulated tones (Parthasarathy & Bartlett, 2011), and reduced post-onset suppression in auditory cortex (Hughes, Turner, Parrish, & Caspary, 2010). Age-related declines in cognitive function can further exacerbate the effects of reduced temporal processing on speech perception, especially in unfavorable listening environments that may mask redundant speech cues (Pichora-Fuller, Schneider, & Daneman, 1995). These studies suggest that training that targets both perceptual and cognitive function is likely to be most effective for addressing age-related concerns.

Mahncke, Bronstone, and Merzenich (2006) have proposed that perceptual training can enhance cognitive performance. In fact, Berry et al. (2010) found that visual discrimination gains were associated with improvement on an untrained working memory task and that the change in N1 amplitude after training (an evoked cortical measure associated with attention; Coch, Sanders, & Neville, 2005) was highly correlated with the extent of working memory improvement. This link between perceptual training and cognitive performance is especially important, because reduced auditory input can impede cognitive function, particularly working memory (Pichora-Fuller, Schneider, & Daneman, 1995). Just as perceptual training is associated with cognitive gains, perhaps cognitive training can improve perceptual performance, especially in light of evidence that age-related declines in sensory function can be compensated by higher-level cognitive and attentional processes (reviewed in Grady, 2012; Li & Lindenberger, 2002; Wong et al., 2009). Wong and colleagues (2010) documented this cognitive compensation in older adults using fMRI to evaluate cortical processing of words in noise. They found reduced activity in auditory cortex but increased activity in prefrontal and pre-cuneus regions – areas associated with memory and attention – in older vs. younger adults. Increased activity in cognitive regions was associated with better performance on a word recognition in noise task, but only in older adults, providing physiological evidence for the importance of memory and attention for hearing in noise in older adults. These results are in line with the Decline Compensation Hypothesis, which states that declines in sensory processing are accompanied by recruitment of general cognitive areas (Cabeza et al., 2004). Both the Berry and the Wong studies reinforce the idea that combining perceptual and cognitive training may engender lasting benefits for communication skills.

Another important consideration is generalization. For any program to be adopted and useful, the training must generalize from trained to untrained tasks or stimuli, especially those important for real world skills (as reviewed in Fahle, 2005). Previous evidence supports generalization for both perceptual

and cognitive tasks in older adults. One study that addressed perceptual generalization showed that although older adults require more sessions to improve performance on a spectro-modulation detection task than younger adults, the generalization of their performance to untrained stimuli was actually better than the generalization in younger adults (Sabin et al., 2013). In a demonstration of transfer to off-task cognitive abilities, older adults who were trained using an auditory-based cognitive software program also had gains on an untrained memory test (Smith et al., 2009). Taken together, these studies demonstrate the potential for cognitive and perceptual training benefits to transfer to more general perceptual and cognitive skills in older adults.

The maintenance of training gains should also be considered when evaluating treatment efficacy, and there is evidence for the maintenance of cognitive gains in older adults. In older participants (i.e., 65–75 years), gains on untrained tasks involving fluid intelligence (e.g., pattern recognition) and speed of processing, but not working memory, were maintained for eight months (Borella, Carretti, Riboldi, & De Beni, 2010). In individuals of more advanced age (i.e., 75–86 years), however, performance on trained tasks was maintained for eight months but the generalization effects were less robust (Borella, Carretti, Zanoni, Zavagnin, & De Beni, 2013). Although no evidence as yet demonstrates the maintenance of perceptual gains in older adults, young adults maintain gains in sentence recognition in noise and enhanced subcortical encoding of a speech syllable in noise for up to six months after using a 20-day speech-in-noise training program (Song, Skoe, Banai, & Kraus, 2012). Given this limited information, there is clearly a need for further investigation into the long-term maintenance of training, especially in older adults. In this study, we investigated the potential maintenance of neural, perceptual, and cognitive training benefits after six months. Based on the Song et al. (2012) findings, we hypothesize that improvements in subcortical temporal processing (decreased latencies and inter-peak variability) and speech-in-noise performance persist for several months. We also hypothesize that gains in speed of processing, but not memory, are maintained after the cessation of training based on Borella et al. (2010).

2. Methods

2.1. Participants

We invited the 67 participants from our original training study to return for a six-month follow-up visit, and 62 participants (34 female) ranging in age from 55 to 70 years returned. We report statistics from the 62 participants who participated in all three sessions, and so there are slight differences in the means and statistical comparisons at post-test from those reported in Anderson, White-Schwoch, Parbery-Clark, and Kraus (2013b). No participants had a history of neurologic conditions, and all participants had normal IQs (≥ 85 on the WASI (Zhu & Garcia, 1999)).

Audiometric thresholds were measured at octave intervals from 0.125 to 8 kHz, including interoctave intervals at 3 and 6 kHz, plus 10 kHz and 12.5 kHz. No changes in threshold (≥ 10 dB) in any participant were noted from the previous sessions. Click-evoked auditory brainstem responses were obtained bilaterally (100 μ s click presented at 80 dB SPL at 31.25 Hz), and no Wave V latency changes (≥ 0.05 ms) from the previous sessions were found in any participant.

All procedures were reviewed and approved by the Northwestern University Institutional Review Board. Participants provided informed consent and were compensated for their time.

2.2. Participant groups

In the original study, participants were randomly assigned to one of two treatment groups, *auditory training* or *active control* (towards the end of the study, targeted enrollment was done to ensure that the groups were matched on age, sex, hearing, and IQ). Both groups completed eight weeks of in-home auditory-based activities on home computers, one hour per day, five days per week, and both groups used headphones (Koss UR/29, Koss Corporation, Milwaukee, WI) to listen

to the training. Of the original 35 members of the *auditory training* group, 30 participated in the final testing session, whereas all 32 members of the *active control* returned. See Fig. 1 for a schematic of the experimental design. The final groups were matched on age, sex, hearing (each frequency 125–12.5 kHz bilaterally), IQ, and test–retest interval (all p 's > 0.10) (Table 1).

2.2.1. Auditory training group

The *auditory training* participants completed 40 h of the Brain Fitness program for 8 weeks (Posit Science, San Francisco, CA), consisting of six modules designed to increase the speed and accuracy of auditory processing: (1) time-order judgment of frequency-modulated sweeps, (2) discrimination between pairs of confusable syllables, (3) recognizing sequences of confusable syllables and words, (4) matching pairs of confusable syllables and words, (5) implementing sequences of commands, and (6) answering questions from stories. The participant's speed of processing is challenged by decreasing the inter-stimulus interval in module 1 with improved performance and by adaptively expanding and contracting the rapidly changing and perceptually vulnerable formant transition of the speech stimuli used in modules 2–6 based on performance. Similarly, memory demands increased (remembering more words, phrases, content) with correct performance. We hypothesized that increased memory demands lead to more focused attention on the CV differences in the training stimuli and engage top-down mechanisms to strengthen sensory input. See Smith et al. (2009) for further details about the training program. Participants received daily online feedback on their performance, and completion of training was verified through online data logging.

2.2.2. Control group

The *active control* group also completed 40 h of in-home computer-based auditory activities for 8 weeks. Participants watched educational DVDs on topics including art, history, and literature and answered questions about the content, and the training was non-adaptive. Only the data of the participants who scored above chance on the questions were included. One participant's scores were below chance level and therefore these data were excluded – indicated in Fig. 1 as “Investigator terminated.”

2.3. Electrophysiology

Electrophysiological methods are identical to those described in Anderson et al. (2013b).

2.3.1. Stimulus

A 170-ms, six-formant speech syllable [da], synthesized in a Klatt-based synthesizer (Klatt, 1980) at a 20 kHz sampling rate was presented in two conditions: in quiet and in two-talker babble background noise (presented at a +10 dB SNR, henceforth referred to as “noise”). After an initial 5 ms stop burst, the stimulus maintained a constant fundamental frequency (F_0) of 100 Hz, with peaks occurring every 10 ms. The lower 3 formants shifted (F_1 : 400→720 Hz; F_2 : 1700→1240 Hz; F_3 : 2580→2500 Hz) during the 50 ms transition from the /d/ to the /a/ but stabilized for the 120 ms steady-state vowel portion. The fourth through sixth formants (F_4 – F_6) remained constant over 170 ms at 3300, 3750, and 4900 Hz, respectively. For participants with hearing loss (thresholds > 20 dB HL at any frequency from 0.25 to 6 kHz for each ear), the frequencies in the [da] were

selectively amplified with the NAL-R algorithm (Byrne & Dillon, 1986) using custom routines in MATLAB (The Mathworks, Inc., Natick, MA) to create customized binaural stimuli to minimize the effects of hearing loss on the responses (Anderson, Parbery-Clark, White-Schwoch, & Kraus, 2013). Seventeen participants in the *auditory training* group and 23 participants in the *active control* received altered stimuli. As hearing thresholds did not change in any participant during the course of the study, each participant received the same stimulus for every testing session.

2.3.2. Recording

Stimuli were presented binaurally in alternating polarities through electrically-shielded insert earphones (ER-3; Etymotic Research, Elk Grove Village, IL) at 80 dB SPL, with an 83 ms interstimulus interval using Neuroscan Stim² (Compumedics, Inc., Charlotte, NC). Subcortical responses were recorded with Ag–AgCl electrodes in a vertical electrode montage (Cz active, forehead ground, earlobe references) with all impedances < 5 k Ω with a 20 kHz sampling rate using Neuroscan Acquire 4.3.

2.3.3. Data reduction

Brainstem responses were offline bandpass filtered from 70 to 2000 Hz in Neuroscan Edit (12 dB/octave roll-off, zero phase-shift), and were epoched with a –40 to 213 ms time window relative to stimulus onset at 0 ms. Responses to the two polarities were added to minimize contamination from stimulus artifact and cochlear microphonic on the response (Campbell, Kerlin, Bishop, & Miller, 2012). Responses were amplitude-baseline corrected relative to the prestimulus period (–40 to 0 ms). Final averages consisted of 6000 artifact-free responses (3000 in each polarity) in each condition.

2.3.4. Data analysis

2.3.4.1. *Peak picking.* Two trained peak-pickers, blind to participant group (*auditory training/active control*) and test session – pre-training (pre), post-training (post), and at six months following cessation of training (post6) – manually identified the major

Table 1

Groups were matched on all demographic data. Means (S.D.s) are displayed for age, sex distributions, test–retest delay (weeks), pure-tone average (.5–4 kHz; dB HL), click wave V latency (ms), IQ (standard score) and MOCA score. Means represent performance at the pre-test.

	Auditory training N=30	Active control N=32
Age (years)	62.3 (3.4)	63.6 (4.1)
Females:Males	16:14	18:14
Test-retest delay in weeks:	9.54 (1.21)	9.9 (2.3)
pre-post		
post-post6	27.3 (3.6)	27.4 (1.6)
Hearing (pure-tone average .5–4 kHz; dB HL)	17.68 (9.8)	18.2 (6.7)
Click wave V latency (ms)	6.0 (0.3)	6.0 (0.4)
IQ (WASI; standard score)	119 (11)	120 (13)




Fig. 1. Flow of participants randomly assigned to *auditory training* or *active control* groups, with a total enrollment of 104 and a final number of 79 after participants were excluded for cognitive, hearing, or neurological reasons. The number of participants who returned for follow-up visits at 8 weeks was 67 and the number at 6 months was 62.

negative-going peaks of interest in the brainstem response. An additional third peak-picker confirmed selections. Peaks of interest in the formant transition of the brainstem response occurred at approximately 34, 44, 54, and 64 ms and in the steady state region at approximately 74, 84, 94, ..., 164 ms. Variability in brainstem peak latencies was determined by calculating the standard deviation of latency differences between adjacent peaks in the frequency following response (FFR) (34–44, 44–54, ..., 154–164) pre- and post-training.

2.3.4.2. Quiet-to-noise correlations. We used a cross-correlation technique (Skoie & Kraus, 2010) to evaluate the degree of response timing shift induced by the babble noise, because masking delays auditory brainstem responses (Anderson, Skoie, Chandrasekaran, & Kraus, 2010; Burkard & Hecox, 1987). This technique shifts the response waveform in noise relative to that in quiet (± 2 ms) until the maximum correlation coefficient (Pearson's r) is found. Fisher z -scores are used for statistical analyses (Cohen & Cohen, 1975). We measured this shift (lag) between the two responses before and after training in the region corresponding to the onset and transition. We expected that the training would have greater effects on the responses in noise and would therefore result in a reduced peak lag in noise relative to a response in quiet. Because this lag is automatically calculated with MATLAB software, this measure serves as an objective confirmation of decreased peak latencies determined through manual peak-picking.

2.4. Behavioral measures

Behavioral methods are identical to those described in Anderson et al. (2013b).

2.4.1. Speech-in-noise recognition

Participants' sentence recognition in noise (SIN) was evaluated with the Quick Speech-in-Noise Test (QuickSIN, Etymotic Research), a non-adaptive clinical measure. Four sets of six sentences are presented in a background of four-talker babble (three female, one male) binaurally at 70 dB HL through insert earphones (ER-2, Etymotic Research). The first sentence in each set is presented at a +25 dB SNR, and the SNR of each of the five subsequent sentences decreases by 5 dB down to a 0 dB SNR. Each sentence contains 5 target words; the total number of correctly repeated target words in each set of six sentences (maximum 30) is subtracted from 25.5 to obtain the dB SNR loss, defined as the difference between the individual's speech-in-noise (SIN) threshold and the average SIN threshold (Killian, Niquette, Gudmundsen, Revit, & Banerjee, 2004). A composite SNR score was obtained by averaging the SNR scores from 4 lists. Different sentence sets were presented at each test session. Lower scores reflect better performance on the task.

2.4.2. Auditory short-term memory

The Woodcock-Johnson III Tests of Cognitive Abilities (Woodcock, McGrew, & Mather, 2001) Memory for Words sub-test was used to evaluate short-term memory. Participants repeat aurally-presented sequences of up to 7 words in the same order as they were presented. Because hearing loss can affect older adults' performance on memory tasks (Baldwin & Ash, 2011), the stimuli were presented with a CD recording and each participant adjusted the volume on the CD player to a comfortable volume. Age-normed scores were used for analysis.

2.4.3. Processing speed

The Woodcock-Johnson III Tests of Cognitive Abilities Visual Matching sub-test was used to evaluate speed of cognitive processing. Participants are presented with 60 printed sets of 6 numbers, ranging from single to triple digits, and in 3 min have to identify (and circle) two identical numbers within each set. Age-normed scores were used for analysis.

2.5. Statistical analyses

Separate repeated measures analyses of variance (RmANOVA) were used to compare neural (inter-peak variability and quiet-to-noise timing) and behavioral measures (speech recognition in noise, short-term memory, and processing speed) across three testing sessions (within subject factor: pre/post/post6) and between groups (between subject factor: *auditory training/active control*). Multivariate RmANOVA were used to compare overall timing changes in the transition (34, 44, 54, 64 ms) and steady state (74, 84, 94, 104, ..., 164 ms) peaks of the brainstem responses. Pearson's correlations were calculated for the *auditory training* group to evaluate relationships among changes in behavioral and neurophysiological variables from the first to the final visit. Normality of all variables was ensured with the Shapiro-Wilk test (all $p > 0.1$) and homogeneity of variance with Levene's test. Mauchly's test of sphericity was used in RmANOVAs, and the Greenhouse-Geisser correction was applied when the sphericity assumption was violated. Bonferroni corrections were applied for follow-up tests in the RmANOVA; p -values reflect two-tailed tests.

3. Results

3.1. Neurophysiology

3.1.1. Latency

In the original study, we found that training improved the timing of neural responses, in particular, the peak latencies in the FFR to a speech syllable. Here we find that the most robust effect on latency in our original study, improved timing in response to the consonant-vowel (CV) transition of a speech syllable presented in noise, persisted for six months. For the peak latency analysis, there was a significant group \times session interaction ($F(2, 61)=2.806, p=0.011$), reflecting a reduction in latency that persisted across the three sessions in the *auditory training* group ($F(1, 29)=2.618, p=0.012$) but not in the *active control* group ($F(1, 31)=1.329, p=0.236$). Post-hoc tests demonstrated a significant reduction in latency from pre to post ($F(1, 29)=6.550, p=0.001$) and pre to post6 ($F(1, 29)=2.807, p=0.046$) that was maintained with no changes from post to post6 ($F(1, 29)=0.607, p=0.661$). There were no latency changes from either pre to post, pre to post6, or from post to post6 in the *active control* group (all p 's > 0.1). The stimulus and grand average responses obtained at the six-month post session in quiet and noise are presented in Fig. 2. The periodicity of the stimulus [da] is seen in the response, with peaks occurring every 10 ms, corresponding to the F_0 of 100 Hz. Latency changes are presented in Fig. 3.

The original results showed a significant, but weaker, reduction in peak latencies in response to the steady-state region of the speech syllable in noise than was found in the transition region, but this training effect did not persist for 6 months – there was no group \times session interaction across the three sessions ($F(2, 61)=1.542, p=0.152$). Table 2 displays mean peak latencies in the noise condition for each group across the three test sessions.

The original study also reported a latency reduction in response to the CV transition of a speech syllable presented in quiet, but this change of latency in quiet was not maintained after six months – there was no group \times test session interaction ($F(2, 61)=1.563, p=0.160$). There had been no training effect in the original study in response to the steady state region in quiet, and similarly no changes were noted at 6 months (i.e., there was no group \times test session interaction) – ($F(2, 61)=1.007, p=0.474$).

3.1.2. Inter-peak variability

In addition to earlier peak latencies in response to speech in noise, the initial study found that training reduced inter-peak variability in the neural responses. The reduction in inter-peak variability in the noise condition was maintained at 6 months during transition and steady-state periods. There was a group \times session interaction ($F(2, 61)=5.860, p=0.005$), with long term training effects across the three sessions in the *auditory training* group ($F(2, 29)=3.622, p=0.040$) but not in the *active control* group ($F(2, 31)=2.219, p=0.126$). Post-hoc tests demonstrated a significant reduction in variability from pre to post ($F(1, 29)=9.979, p=0.004$) and pre to post6 ($F(1, 29)=4.391, p=0.047$) with no significant changes from post to post6 sessions in the *auditory training* group ($F(1, 29)=0.13, p=0.910$). There were no changes from either pre to post, pre to post6 or from post to post6 in the *active control* group (all p 's > 0.1).

3.1.3. Quiet-to-noise timing

In the original study we also observed a reduction in quiet-to-noise timing shift, or the extent to which the babble noise delayed response peaks. This reduction was not maintained – group \times session interaction and paired t -tests all resulted in p 's > 0.1 .

3.2. Behavioral

In our original study, we observed behavioral improvements in speed of processing, short-term memory, and speech-in-noise recognition. Group means and S.D.s for pre- and post-training and post6 behavioral measures are displayed in Table 3. There were no significant differences in *auditory training* and *active control* group performance on these measures at pre-test (SIN recognition: $t(65)=0.501$, $p=0.618$; short-term memory: $t(65)=1.422$, $p=0.160$; processing speed: $t(65)=0.678$, $p=0.500$).

The improvement in speed of processing was maintained at the six-month follow-up session in the *auditory training* group as evidenced by a group \times session interaction ($F(2, 61)=3.192$, $p=0.048$) with a long-term training effect across the three test sessions in the *auditory training* group ($F(2, 29)=4.317$, $p=0.023$) but not in the *active control* group ($F(2, 31)=1.730$, $p=0.195$). Post-hoc testing indicated that for the *auditory training* group, there was improvement in speed of processing from pre to post ($F(1, 29)=6.556$, $p=0.016$) and pre to post6 ($F(1, 29)=6.940$, $p=0.014$) which was maintained from post to post6 ($F(1, 29)=0.043$, $p=0.837$). In the *active control* group, there were no changes

in speed of processing from pre to post or from pre to post6 (all p 's > 0.1).

However, improvements in speech-in-noise performance and in short-term memory were not retained. There was no group \times session interaction for speech-in-noise performance ($F(2, 61)=0.613$, $p=0.545$). There was a significant group \times session interaction for memory ($F(2, 61)=5.292$, $p=0.008$), because the training effect seen in the *auditory training* group at the post session ($F(1, 29)=8.282$, $p < 0.001$) was not maintained at post6 ($F(1, 29)=3.937$, $p=0.057$), and there were no changes seen from pre to post or post to post6 in the *active control* group (all p 's > 0.1). Changes in behavioral performance on speech-in-noise recognition, short-term memory and speed of processing between pre and post and between pre and post6 for both groups are depicted in Fig. 4.

3.3. Brain-Behavior Correlations

Using data from the *auditory training* group, we found that improvements in processing speed were related to decreases in peak variability from pre to post6 (Fig. 5). In addition, decreases in peak variability were related to decreases in transition peak




Fig. 2. The periodicity of the evoking 170-ms stimulus [da] with a fundamental frequency of 100 Hz is represented in the grand average response obtained at the six-months post visit to the stimulus both in quiet (gray) and in noise (black), with peaks occurring at a frequency of every 10 ms.




Fig. 3. In noise, the reduction in peak latencies in the region corresponding to the consonant–vowel transition (20–60 ms, marked with brackets) that was found after 8 weeks of training was maintained at the six-month follow-up visit. To facilitate visualization of the data across time, peak latency values are normalized by subtracting individual absolute latencies from the expected response latencies for each peak based on average data in young adults, taking into account stimulus characteristics and neural lag (Anderson et al., 2012). Therefore, faster timing corresponds to smaller differences between actual and expected latency values. Expected latencies were 34, 44, 54, 64, etc., until 164 ms for peaks during the transition and steady-state. No latency changes were seen in the *active control* group. * $p < 0.05$ – significance value for the pre to post-6 change in latency. Error bars = ± 1 S.E.

Table 2

Means (S.E.s) are displayed for peak latencies occurring every 10 ms from 34 to 164 ms in response to the syllable [da] presented in two-talker babble obtained during the pre-, post-8 weeks, and post-6 months sessions for each group.

Peak latencies (ms)	Auditory training			Active control		
	Pre	Post	Post 6	Pre	Post	Post 6
34	35.61 (0.22)	34.90 (0.21)	35.17 (0.18)	35.22 (0.22)	35.24 (0.20)	34.95 (0.16)
44	45.02 (0.23)	44.23 (0.17)	44.29 (0.18)	46.24 (0.15)	46.19 (0.20)	45.96 (0.13)
54	54.80 (0.17)	54.41 (0.13)	54.46 (0.13)	54.47 (0.15)	54.45 (0.15)	54.34 (0.12)
64	64.88 (0.16)	64.52 (0.13)	64.38 (0.10)	64.49 (0.12)	64.62 (0.18)	64.52 (0.12)
74	74.78 (0.11)	74.53 (0.08)	74.55 (0.10)	74.63 (0.14)	74.67 (0.18)	74.46 (0.13)
84	84.87 (0.20)	84.51 (0.09)	84.48 (0.08)	84.64 (0.16)	84.67 (0.18)	84.44 (0.09)
94	94.74 (0.12)	94.35 (0.09)	94.62 (0.10)	94.60 (0.14)	94.60 (0.17)	94.43 (0.09)
104	104.81 (0.17)	104.48 (0.09)	104.64 (0.10)	104.50 (0.09)	104.58 (0.16)	104.42 (0.09)
114	114.71 (0.08)	114.48 (0.08)	114.68 (0.13)	114.58 (0.13)	114.65 (0.21)	114.41 (0.08)
124	124.78 (0.12)	124.50 (0.07)	124.67 (0.12)	124.44 (0.08)	124.67 (0.21)	124.44 (0.09)
134	134.77 (0.13)	134.52 (0.08)	134.67 (0.11)	134.58 (0.13)	134.58 (0.13)	134.47 (0.09)
144	144.80 (0.14)	144.55 (0.07)	144.62 (0.08)	144.44 (0.09)	144.71 (0.22)	144.38 (0.06)
154	154.84 (0.23)	154.46 (0.09)	154.53 (0.09)	154.65 (0.13)	154.58 (0.16)	154.42 (0.09)
164	164.77 (0.15)	164.45 (0.07)	164.52 (0.09)	164.51 (0.09)	164.70 (0.22)	164.43 (0.09)

Table 3

Pre-, post-8 week, and post-6 month behavioral means (S.E.s) for the *auditory training* and *active control* groups. The speed of processing and short-term memory means are age-normed standard scores. The speech-in-noise (SIN) score means are in dB SNR and represent the difference between the individual's SIN threshold and the average SIN threshold – the SNR loss (Killion et al., 2004).

	Auditory training			Active control		
	Pre	Post	Post 6	Pre	Post	Post 6
Speed of processing (standard score)	107.93 (2.15)	112.13 (2.53)	112.57 (2.61)	109.74 (1.86)	107.33 (1.85)	110.71 (2.15)
Speech-in-noise (dB SNR loss)	1.08 (0.21)	0.36 (0.25)	0.81 (0.19)	1.06 (0.17)	0.65 (0.21)	0.84 (0.20)
Short-term memory (standard score)	106.90 (2.88)	114.87 (2.35)	110.80 (2.73)	110.00 (1.70)	110.85 (2.13)	110.68 (2.21)

latency. There were no other significant correlations among the speech-in-noise and short-term memory measures and neurophysiological measures from pre to post6. In addition, there were no significant correlations among any of the variables from pre to post or from post to post6. See Table 4 for correlation values for pre to post6 changes in behavioral and neurophysiological measures in the *auditory training* group.

4. Discussion

The maintenance of the training benefits produced by commercial software programs is an important index of training success. Here we report auditory training effects on subcortical responses to speech and behavioral measures of speed of processing, speech-in-noise recognition, and short-term memory in older adults, some of which persisted for six months. We found that the most robust effects in the original training study, reductions in peak latencies in response to the CV transition of a speech syllable and interpeak variability, were maintained for six months after the completion of training. Interestingly, improvement in our behavioral measure of speed of processing also persisted for six months, whereas memory and speech-in-noise performance gains did not. See Table 5 for a summary of the initial training effects and their maintenance after six months.

4.1. Maintenance of changes in response peak timing

To add to previous reports of cortical auditory plasticity in humans (Du et al., 2013; Hayes, Warrier, Nicol, Zecker, & Kraus, 2003; Tremblay, Shahin, Picton, & Ross, 2009), mounting evidence now suggests that short-term training can modify subcortical auditory processing (Carcagno & Plack, 2011; Chandrasekaran,

Kraus, & Wong, 2012; de Boer & Thornton, 2008; Song et al., 2012). One of these studies specifically examined the maintenance of these effects in young adults (Song et al., 2012). Although the current study is the first to examine maintenance of neural changes to short-term training initiated by older adults, the benefits of music lessons received during childhood can persist into older adulthood (White-Schwoch, Carr, Anderson, Strait, & Kraus, 2013). In the White-Schwoch et al. study, older adults with music training in childhood had earlier subcortical peak latencies in response to speech than those who had no music training, with years of training correlating with peak latencies. The authors proposed that this training early in life may prime the auditory system to selectively respond to auditory inputs to better encode the acoustic complexity and the salient information of the signal. The results from the current study suggest that short-term training initiated later in life can instill changes that persist for up to six months.

The neural timing benefits (earlier peak timing) in the current study and that of White-Schwoch et al. (2013) served to offset one of the most pervasive aspects of age-related declines – an overall slowing in neural speed of processing and a decrease in synchronous responses. In our original study, we speculated that an increase in inhibitory neurotransmission in auditory midbrain occurred in response to the training, potentially due to training demands of attention to rapidly changing details of the CV transition. This conjecture was based in part on animal models of auditory plasticity that have demonstrated increased numbers of inhibitory gamma-aminobutyric acid (GABA) receptors in auditory cortex following auditory training (Guo et al., 2012) and that pharmacologic treatment with a GABAergic agent can improve gap detection performance (Gleich, Hamann, Klump, Kittel, & Strutz, 2003). GABA is necessary for the precise temporal coding of the fast-changing formants in consonant–vowel transitions (as reviewed in Caspary, Ling, Turner, & Hughes, 2008).




Fig. 4. In the *auditory training* group (red circles), the increase in processing speed (standard score) seen at the post-training visit persisted for 6 months (A), but the training-induced enhancements in short-term memory (standard score) (B); and sentence recognition in noise (dB SNR loss – lower scores indicate better performance) (C) did not persist. There were no initial changes in processing speed or memory in the *active control* group (blue squares), and the initial change in speech-in-noise recognition did not persist. * $p < 0.05$ – significance value for the group \times session interaction. Error bars = ± 1 S.E. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 5. Changes in processing speed relate to changes in brainstem peak variability from pre to post6 in the *auditory training* group, such that individuals who experienced greater improvement in processing speed were more likely to have reduced peak variability.

Table 4

Correlation values are listed for pre to post6 change measures in peak variability, transition latency (peak 44), speed of processing, short-term memory, and speech-in-noise recognition in the *auditory training* group. Relationships between changes in speed of processing and peak variability and between peak variability and transition latency were significant.

	Processing speed	Short-term memory	Speech-in-noise	Peak variability	Transition peak latency
Processing speed	1	-0.160	-0.070	-0.499**	-0.126
Short-term memory	-0.160	1	0.110	-0.035	0.045
Speech-in-noise	-0.070	0.110	1	-0.160	-0.225
Peak variability	-0.499**	-0.035	-0.160	1	0.575**
Transition peak latency	-0.126	0.045	-0.225	0.575**	1

** $p < 0.01$.

Table 5

Summary of training (pre to post) and maintenance effects (pre to post6) in the *auditory training* group with asterisks indicating significance levels for each effect. The neurophysiology effects that were maintained were the reduction in peak latencies to the CV transition presented in two-talker babble and the reduction in inter-peak variability. The improvement in speed of processing was the only behavioral effect that persisted.

Training effect	Pre to post	Pre to post-6
<i>Neurophysiology</i>		
Reduction in latency in quiet		
Transition	*	
Steady state		
Reduction in latency in noise		
Transition	***	*
Steady state	*	
Reduction in inter-peak variability	*	*
Reduction in quiet-to-noise timing shift	*	
<i>Behavior</i>		
Speech-in-noise perception	**	
Short-term memory	**	
Speed of processing	**	*

* $p < 0.05$.
 ** $p < 0.01$.
 *** $p < 0.001$.

Cholinergic input to auditory cortex from nucleus basalis in the basal forebrain is required to induce plasticity in auditory perception (Bakin & Weinberger, 1996; Leach, Nodal, Cordery, King, & Bajo, 2013; Puckett, Pandya, Moucha, Dai, & Kilgard, 2007). GABA neurons in the mesolimbic reward system (ventral tegmental area and nucleus accumbens) target cholinergic interneurons to modulate learning (Creed, Ntamati, & Tan, 2014). The training used in our study, with its intrinsic rewards from feedback on improved performance, may have resulted in changes in these neurotransmitter levels, which may in turn have contributed to a reduction in neural variability (Zinke et al., 2006). Due to the extensive series of descending corticofugal projections, we suggest that this GABAergic increase propagated from auditory cortex to midbrain, leading to more precise (i.e., more synchronous and faster) temporal coding of fast-changing sounds, eventually creating long-lasting changes as reflected in our studies. Assuming that the initial changes in peak latencies arose from an increase in inhibitory neurotransmission, we speculate that these changes persisted to some extent for six months. In our original study, the greatest latency changes were found in response to the CV transition presented in noise, and these were the only latency changes that persisted for six months. These results are consistent with our neurotransmitter view, because inhibitory function is especially important for coding rapidly changing sounds (Casparly et al., 2008), as found in the CV transition.

4.2. Maintenance of changes in cognitive skills

Gains in speed of processing, but not in short-term memory, persisted for six months. Interestingly, reduction in peak variability related to improvements in processing speed over the course of the study, while changes in sentence recognition in noise and short-term memory did not relate to any neurophysiologic changes. The relationship between changes in processing speed and peak variability suggests that peak variability in neural responses to sensory inputs may be a factor in processing speed deficits in older adults.

The persistence of benefits for speed of processing, but not for memory, are consistent with the findings of Borella et al. (2010), who showed that older adults maintained training enhancements for fluid intelligence and speed of processing but not for memory. In their study they trained verbal working memory using an adaptive procedure and then determined whether this training resulted in performance gains on the trained task (following 2 weeks of training), short-term transfer effects, and long-term maintenance (8 months) of gains. Salthouse (1996) proposed that speed of processing is a basic aspect of cognitive function that declines with age and underlies changes in specific cognitive skills such as working memory. Borella et al. postulated that the training causes a boost in overall cognitive processing (i.e. speed of processing) but that the change for specific skills, such as working memory, is not robust enough to resist erosion over time. In our training regimen, the reduction in inter-stimulus intervals in module 1 or compression of CV transitions in modules 2–6 (see Section 2.2.1) may have necessitated an increase in processing speed that continued into everyday tasks and was therefore maintained for an additional six months. Of course, there are also opportunities to practice memory in everyday life. However, a higher practice threshold may be necessary to maintain improved memory skills, and this maintenance may require more initial training or explicit practice.

Aging may also affect retention of enhanced memory performance. Li et al. (2008) compared working memory training effects in younger vs. older adults. Both age groups improved on the trained task and performance generalized to a more demanding working memory task. This performance improvement was maintained for three months in the young adults; the older adults showed a decline relative to post-training session, but their end performance at three months was still significantly better than at pre-test. Another study found maintenance in working memory gains for 18 months in both younger and older adults; however, in the older adults there was no transfer of learning to an untrained task (Dahlin, Nyberg, Backman, & Neely, 2008).

4.3. Implications for speech-in-noise perception

We were surprised that the improvement in SIN recognition did not hold, especially given the likelihood of having many opportunities to practice this skill in real-world environments after training had ceased. Older adults, both with normal hearing and with hearing loss, have deficits in auditory temporal processing that lead to impaired perception. For example, older adults require longer consonant transition durations to discriminate between words that differ only on this temporal cue (e.g., “beat” vs. “wheat”) in isolated words (Gordon-Salant, Yeni-Komshian, Fitzgibbons, & Barrett, 2006) and sentences (Fitzgibbons, Gordon-Salant, & Friedman, 2006). Furthermore, older adults have more difficulty recognizing time-compressed speech materials presented in quiet (Gordon-Salant & Fitzgibbons, 2001) and at different SNRs (Jafari, Omidvar, & Jafarloo, 2013), suggesting that older adults have difficulty processing brief temporal cues that are characteristic of rapid speech. Therefore, given that the subcortical

temporal processing gains in our study were maintained, we would have expected to see similar results for our perceptual measures. We suspect that factors other than enhanced neural timing were at least partly responsible for the initial gains in speech perception. The QuickSIN sentences used to assess SIN recognition are long – 8–11 words – and therefore impose strong demands on short-term memory and attention. While we believe that the both cognitive and sensory processing factors contribute to speech-in-noise performance (Anderson et al., 2013a), the fact that improvement in QuickSIN scores did not persist despite the maintenance of improvements in subcortical peak timing and variability has led us to believe that the initial improvements in QuickSIN may have been driven in part by the cognitive components of the training. The training did not explicitly teach participants strategies for improving their ability to hear in noise. Perhaps, actual listening-in-noise exercises may be necessary to engender lasting changes in skills or strategies used while listening to speech in adverse listening environments. In our literature search, we found only a few reports of long-term maintenance of auditory training. The benefits of digits-in-noise training in older individuals with cochlear implants (ages 46–78) were maintained for one month after cessation of training and generalized to untrained tests of speech-in-noise performance, but no results were reported for a longer period of maintenance (Oba, Fu, & Galvin, 2011). Words-in-noise training produced significant improvement in older listeners (ages 65–75) that were maintained six months after the cessation of training, but the training gains did not generalize to recognition of the trained words in sentences (Burk, Humes, Amos, & Strauser, 2006). Listening and Communication Enhancement (LACE™, Neurotone, Redwood City, CA) is an example of a home-based computerized training program that explicitly trains listening in competing or background noise, in addition to other top-down strategies. Young adults who underwent LACE training showed maintenance effects lasting six months that generalized to two non-trained measures of sentence recognition in noise (Song et al., 2012). This study also showed a maintenance of enhanced subcortical responses to speech, with greater representation of the fundamental frequency in response to a speech syllable embedded in noise after 4 weeks of training with LACE, and then again at 6 months after the discontinuation of LACE training.

One explanation for the difference in maintenance outcomes between our study and the Song et al. (2012) study may be the study populations: Song et al. trained young adults whereas we trained older adults. Training-induced changes in young adults may be more robust and resistant to change than those in older adults. In fact, two studies that trained perception of time-compressed speech found that younger, but not older, adults retained training benefits (Barcroft et al., 2011; Peelle & Wingfield, 2005).

Younger adults may be more efficient in the engagement of the neural circuits necessary to learn a new skill compared to older adults, and this enhanced efficiency may account for the persistence of training benefits in younger adults. Park and Reuter-Lorenz (2009) introduced the term “scaffolding” to describe the engagement of complementary and redundant neural circuits in response to challenges that might be posed during a learning task or when sensory input is reduced. Although older adults use scaffolding as a compensatory mechanism, their scaffolding is less efficient than that in young adults; therefore, training-induced changes may not persist in the older population.

4.4. Ingredients for successful learning

Although memory exercises may be insufficient in and of themselves to effect lasting changes in SIN recognition, memory

does play an important role in successful listening in noise in both younger and older adults (Pichora-Fuller et al., 1995). In fact, memory may be more important for older listeners than younger listeners, an idea that would be broadly consistent with the Cognitive Compensation Hypothesis, which suggests that cognitive processes, such as memory and attention, compensate for sensory losses in older adults (Wong, Ettlenger, Sheppard, Gunasekera, & Dhar, 2010). Even in young listeners, cognitive influences can compensate for degraded sensory input (Nahum, Nelken, & Ahissar, 2008).

In older adults, neuroimaging studies support the idea that cortical areas dedicated to executive function have greater activation during speech-in-noise listening tasks in older adults than in younger adults (Wong et al., 2010). Furthermore, a structural equation model of speech-in-noise perception has demonstrated a strong role for both cognitive factors and central auditory processing in understanding speech in noise, with nonsignificant contributions to the model from peripheral function (Anderson et al., 2013a). Therefore, cognitive training (e.g., memory, attention) may be an important component of SIN training, but the training dosage may need to be increased or booster sessions may be necessary to maintain learning. Older adults who received supplemental booster sessions in cognitive training on either episodic memory, reasoning, or speed of processing tasks had better maintenance of training gains in reasoning and speed of processing, but not in memory, 11 months later when compared to groups who had received similar training without the booster sessions (Ball et al., 2002). The training program we used did not include the option of booster sessions following completion of training, but we recognize that this is an important consideration in the selection or design of training programs for any age, particularly in older adults whose performance is less likely to be stable over time compared to that of younger adults. This is an important area of follow-up in future dosage studies.

4.5. Limitations

In this study we wished to establish an effect using a commercially available training program, but one of the limitations of selecting a pre-packaged program is its lack of flexibility. For example, although our primary interest was in the training effects on untrained tasks (rather than in program-trained perceptual tasks), it would have been beneficial to assess performance on these trained tasks after six months. Doing so would have allowed us to determine, among other things, if there is a relationship between behavioral performance on trained and untrained tasks and neurophysiologic measures. This information would be useful in guiding the development of future training programs and in tracking of trainee progress.

We used a visual measure of processing speed, the Visual Memory subtest of the Woodcock–Johnson III Tests of Cognitive Abilities, rather than an auditory measure; therefore, we cannot with certainty conclude these results generalize to the auditory domain. However, the fact that changes in a behavioral measure of visual processing speed correlate with changes in a neurophysiological measure of auditory processing suggests that the training may have had a domain general rather than a domain specific effect. It would be interesting to compare training effects on visual vs. auditory processing speed measures in a future study design.

Finally, as mentioned above, performance on the QuickSIN may have been affected by short-term memory demands. For this reason, it would be beneficial to include speech recognition measures that are less affected by memory, such as the Bamford–Kowal–Bench Speech-in-Noise Test and the Words-in-Noise Test (Wilson, McArdle, & Smith, 2007), in future training studies designed to improved speech-in-noise performance.

4.6. Future directions

Future work should consider factors that are important for the maintenance of training. One such factor is compliance with the training regimen. Programs that are engaging may enhance compliance with training (Nacke, Nacke, & Lindley, 2009). Individuals are less likely to return to booster sessions when the task becomes less interesting. Recent studies have examined the use of engaging video games to improve executive function (Anguera et al., 2013) and to enhance learning capacity (i.e. learning to learn) (Bavelier, Green, Pouget, & Schrater, 2012), as they may improve compliance by providing a more entertaining and rewarding training system. We envision future applications of similar technology for improving speech perception in older adults.

Future work should also consider the role of hearing loss in training outcomes. We did not include any individual with more than a mild-moderate sensorineural hearing loss, and given the complex interactions between working memory, hearing loss, and the temporal coding of speech (Anderson, Parbery-Clark, White-Schwoch, Drehobl, & Kraus, 2013; Lin et al., 2013; Rudner, Rönnerberg, & Lunner, 2011), older adults with more severe hearing loss may have a different training and maintenance profile (Anderson, White-Schwoch, Choi, & Kraus, 2013). We also acknowledge that there may be individual differences in training outcomes independent of hearing status and future work should investigate factors that might contribute to success (or not) in older adults.

4.7. Conclusions

These findings demonstrate that short-term training-induced enhancements in auditory temporal precision persist for at least six months in older adults. These results are particularly important in light of the known pervasive declines in neural temporal precision in older adults (Anderson et al., 2012; Hughes et al., 2010; Lister et al., 2011). However, we did not find equivalent maintenance across all the training effects that we reported in Anderson et al. (2013b). Specifically, while most of the neural changes persisted (earlier peak timing, reduction in peak variability), speed of processing was the only behavioral measure that maintained improvement for six months. The improvements in sentence-in-noise recognition and short-term memory did not persist. The fact that we did not find maintenance across all domains suggests that training may not engender a domain general plasticity but instead may train a series of systems with different parameters and limits for changes. Broadly speaking, our results suggest that short-term training can instill changes in a number of cognitive and sensory functions that have different conditions for improvement and long-term maintenance and may suggest that diverse training regimens are most effective for lasting improvements.

Acknowledgments

The authors thank Sarah Drehobl, Rafael Escobedo, and Samantha O'Connell for their assistance with data collection. We thank Trent Nicol, Laurence Anderson, and three anonymous reviewers for their helpful reviews and suggestions for the manuscript. We also thank our participants for their willingness to return for multiple testing sessions. This work was supported by the National Institutes of Health (T32 DC009399 and R01 DC10016) and the Knowles Hearing Center.

References

- Anderson, S., Parbery-Clark, A., White-Schwoch, T., Dreihobl, S., & Kraus, N. (2013). Effects of hearing loss on the subcortical representation of speech cues. *The Journal of the Acoustical Society of America*, *133*, 3030–3038.
- Anderson, S., Parbery-Clark, A., White-Schwoch, T., & Kraus, N. (2012). Aging affects neural precision of speech encoding. *The Journal of Neuroscience*, *32*, 14156–14164.
- Anderson, S., Parbery-Clark, A., White-Schwoch, T., & Kraus, N. (2013). Auditory brainstem response to complex sounds predicts self-reported speech-in-noise performance. *Journal of Speech, Language & Hearing Research*, *56*, 31–43.
- Anderson, S., Skoe, E., Chandrasekaran, B., & Kraus, N. (2010). Neural timing is linked to speech perception in noise. *The Journal of Neuroscience*, *30*, 4922–4926.
- Anderson, S., White-Schwoch, T., Choi, H. J., & Kraus, N. (2013). Training changes processing of speech cues in older adults with hearing loss. *Frontiers in Systems Neuroscience*, *7*.
- Anderson, S., White-Schwoch, T., Parbery-Clark, A., & Kraus, N. (2013a). A dynamic auditory-cognitive system supports speech-in-noise perception in older adults. *Hearing Research*, *300*, 18–32.
- Anderson, S., White-Schwoch, T., Parbery-Clark, A., & Kraus, N. (2013b). Reversal of age-related neural timing delays with training. *Proceedings of the National Academy of Sciences – USA*, *110*, 4357–4362.
- Anguera, J., Boccanfuso, J., Rintoul, J., Al-Hashimi, O., Faraji, F., Janowich, J., et al. (2013). Video game training enhances cognitive control in older adults. *Nature*, *501*, 97–101.
- Bakin, J. S., & Weinberger, N. M. (1996). Induction of a physiological memory in the cerebral cortex by stimulation of the nucleus basalis. *Proceedings of the National Academy of Sciences – USA*, *93*, 11219–11224.
- Baldwin, C. L., & Ash, I. K. (2011). Impact of sensory acuity on auditory working memory span in young and older adults. *Psychology and Aging*, *26*, 85–91.
- Ball, K., Berch, D. B., Helmers, K. F., Jobe, J. B., Leveck, M. D., Mariske, M., et al. (2002). Effects of cognitive training interventions with older adults: A randomized controlled trial. *Journal of the American Medical Association*, *288*, 2271–2281.
- Barcroft, J., Sommers, M. S., Tye-Murray, N., Mauzé, E., Schroy, C., & Spehar, B. (2011). Tailoring auditory training to patient needs with single and multiple talkers: Transfer-appropriate gains on a four-choice discrimination test. *International Journal of Audiology*, *50*, 802–808.
- Bavelier, D., Green, C. S., Pouget, A., & Schrater, P. (2012). Brain plasticity through the life span: Learning to learn and action video games. *Annual Review of Neuroscience*, *35*, 391–416.
- Berry, A. S., Zanto, T. P., Clapp, W. C., Hardy, J. L., Delahunt, P. B., Mahncke, H. W., et al. (2010). The influence of perceptual training on working memory in older adults. *Plos One*, *5*, e11537.
- Borella, E., Carretti, B., Riboldi, F., & De Beni, R. (2010). Working memory training in older adults: Evidence of transfer and maintenance effects. *Psychology and Aging*, *25*, 767.
- Borella, E., Carretti, B., Zononi, G., Zavagnin, M., & De Beni, R. (2013). Working memory training in old age: An examination of transfer and maintenance effects. *Archives of Clinical Neuropsychology*, *28*, 331–347.
- Burk, M. H., Humes, L. E., Amos, N. E., & Strauser, L. E. (2006). Effect of training on word-recognition performance in noise for young normal-hearing and older hearing-impaired listeners. *Ear & Hearing*, *27*, 263–278.
- Burkard, R., & Hecox, K. E. (1987). The effect of broadband noise on the human brain-stem auditory evoked response. III. Anatomic locus. *The Journal of the Acoustical Society of America*, *81*, 1050.
- Byrne, D., & Dillon, H. (1986). The National Acoustic Laboratories' (NAL) new procedure for selecting the gain and frequency response of a hearing aid. *Ear & Hearing*, *7*, 257–265.
- Cabeza, R., Daselaar, S. M., Dolcos, F., Prince, S. E., Budde, M., & Nyberg, L. (2004). Task-independent and task-specific age effects on brain activity during working memory, visual attention and episodic retrieval. *Cerebral Cortex*, *14*, 364–375.
- Campbell, T., Kerlin, J., Bishop, C., & Miller, L. (2012). Methods to eliminate stimulus transduction artifact from insert earphones during electroencephalography. *Ear and Hearing*, *33*, 144–150.
- Carcagno, S., & Plack, C. (2011). Subcortical plasticity following perceptual learning in a pitch discrimination task. *Journal of the Association for Research in Otolaryngology*, *12*, 89–100.
- Caspary, D. M., Ling, L., Turner, J. G., & Hughes, L. F. (2008). Inhibitory neurotransmission, plasticity and aging in the mammalian central auditory system. *Journal of Experimental Biology*, *211*, 1781–1791.
- Chandrasekaran, B., Kraus, N., & Wong, P. C. M. (2012). Human inferior colliculus activity relates to individual differences in spoken language learning. *Journal of Neurophysiology*, *107*, 1325–1336.
- Clinard, C., & Tremblay, K. (2013). Aging degrades the neural encoding of simple and complex sounds. *Journal of the American Academy of Audiology*, *24*, 590–599.
- Coch, D., Sanders, L. D., & Neville, H. J. (2005). An event-related potential study of selective auditory attention in children and adults. *Journal of Cognitive Neuroscience*, *17*, 605–622.
- Cohen, J., & Cohen, P. (1975). *Applied multiple regression/correlation analysis for the behavioral sciences*. Hillsdale, NJ: New York: Lawrence Erlbaum Associates.
- Creed, M. C., Ntamati, N. R., & Tan, K. R. (2014). VTA GABA neurons modulate specific learning behaviours through the control of dopamine and cholinergic systems. *Frontiers in Behavioral Neuroscience*, *8*, 1–7.
- Dahlin, E., Nyberg, L., Backman, L., & Neely, A. S. (2008). Plasticity of executive functioning in young and older adults: Immediate training gains, transfer, and long-term maintenance. *Psychology and Aging*, *23*, 720.
- de Boer, J., & Thornton, A. R. D. (2008). Neural correlates of perceptual learning in the auditory brainstem: Efferent activity predicts and reflects improvement at a speech-in-noise discrimination task. *The Journal of Neuroscience*, *28*, 4929–4937.
- Du, Y., He, Y., Arnott, S. R., Ross, B., Wu, X., Li, L., et al. (2013). Rapid tuning of auditory “What” and “Where” pathways by training. *Cerebral Cortex*, bht251.
- Fahle, M. (2005). Perceptual learning: Specificity versus generalization. *Current Opinion in Neurobiology*, *15*, 154–160.
- Fitzgibbons, P. J., Gordon-Salant, S., & Friedman, S. A. (2006). Effects of age and sequence presentation rate on temporal order recognition. *The Journal of the Acoustical Society of America*, *120*, 991–999.
- Gleich, O., Hamann, I., Klump, G. M., Kittel, M., & Strutz, J. (2003). Boosting GABA improves impaired auditory temporal resolution in the gerbil. *NeuroReport*, *14*, 1877–1880.
- Gordon-Salant, S., & Fitzgibbons, P. (2001). Sources of age-related recognition difficulty for time-compressed speech. *Journal of Speech, Language, and Hearing Research*, *44*, 709–719.
- Gordon-Salant, S., Yeni-Komshian, G. H., Fitzgibbons, P. J., & Barrett, J. (2006). Age-related differences in identification and discrimination of temporal cues in speech segments. *The Journal of the Acoustical Society of America*, *119*, 2455–2466.
- Grady, C. (2012). The cognitive neuroscience of ageing. *Nature Reviews Neuroscience*, *13*, 491–505.
- Guo, F., Zhang, J., Zhu, X., Cai, R., Zhou, X., & Sun, X. (2012). Auditory discrimination training rescues developmentally degraded directional selectivity and restores mature expression of GABA_A and AMPA receptor subunits in rat auditory cortex. *Behavioural Brain Research*, *229*, 301–307.
- Hayes, E. A., Warrier, C. M., Nicol, T. G., Zecker, S. G., & Kraus, N. (2003). Neural plasticity following auditory training in children with learning problems. *Clinical Neurophysiology*, *114*, 673–684.
- Hughes, L. F., Turner, J. G., Parrish, J. L., & Caspary, D. M. (2010). Processing of broadband stimuli across A1 layers in young and aged rats. *Hearing Research*, *264*, 79–85.
- Jafari, Z., Omidvar, S., & Jafarloo, F. (2013). Effects of ageing on speed and temporal resolution of speech stimuli in older adults. *Medical Journal of the Islamic Republic of Iran (MJIRI)*, *27*, 195–203.
- Killion, M., Niquette, P., Gudmundsen, G., Revit, L., & Banerjee, S. (2004). Development of a quick speech-in-noise test for measuring signal-to-noise ratio loss in normal-hearing and hearing-impaired listeners. *Journal of the Acoustical Society of America*, *116*, 2395–2405.
- Klatt, D. (1980). Software for a cascade/parallel formant synthesizer. *Journal of the Acoustical Society of America*, *67*, 971–995.
- Kraus, N., & Chandrasekaran, B. (2010). Music training for the development of auditory skills. *Nature Reviews Neuroscience*, *11*, 599–605.
- Krishnan, A., Gandour, J. T., & Bidelman, G. M. (2010). The effects of tone language experience on pitch processing in the brainstem. *Journal of Neurolinguistics*, *23*, 81–95.
- Leach, N. D., Nodal, F. R., Cordery, P. M., King, A. J., & Bajo, V. M. (2013). Cortical cholinergic input is required for normal auditory perception and experience-dependent plasticity in adult ferrets. *The Journal of Neuroscience*, *33*, 6659–6671.
- Li, K. Z. H., & Lindenberger, U. (2002). Relations between aging sensory/sensorimotor and cognitive functions. *Neuroscience Biobehavioral Reviews*, *26*, 777–783.
- Li, S.-C., Schmiedek, F., Huxhold, O., Rocke, C., Smith, J., & Lindenberger, U. (2008). Working memory plasticity in old age: Practice gain, transfer, and maintenance. *Psychology and Aging*, *23*, 731.
- Lin, F. R., Yaffe, K., Xia, J., Xue, Q. L., Harris, T. B., & Purchase-Helzner, E. (2013). Hearing loss and cognitive decline in older adults. *Journal of the American Medical Association Internal Medicine*, *173*, 293–299.
- Lister, J. J., Maxfield, N. D., Pitt, G. J., & Gonzalez, V. B. (2011). Auditory evoked response to gaps in noise: Older adults. *International Journal of Audiology*, *50*, 211–225.
- Mahncke, H. W., Bronstone, A., & Merzenich, M. M. (2006). Brain plasticity and functional losses in the aged: Scientific bases for a novel intervention. In: R. M. Aage (Ed.), *Progress in brain research*, Vol. 157 (pp. 81–109). Amsterdam: Elsevier.
- Nacke, L. E., Nacke, A., & Lindley, C. A. (2009). Brain training for silver gamers: Effects of age and game form on effectiveness, efficiency, self-assessment, and gameplay experience. *Cyberpsychology Behavior*, *12*, 493–499.
- Nahum, M., Nelken, I., & Ahissar, M. (2008). Low-level information and high-level perception: The case of speech in noise. *PLoS biology*, *6*, e126.
- Oba, S. I., Fu, Q.-J., & Galvin, J. J. I. (2011). Digit training in noise can improve cochlear implant users' speech understanding in noise. *Ear & Hearing*, *32*, 573–581.
- Park, D. C., & Reuter-Lorenz, P. (2009). The adaptive brain: Aging and neurocognitive scaffolding. *Annual Reviews of Psychology*, *60*, 173.
- Parthasarathy, A., & Bartlett, E. L. (2011). Age-related auditory deficits in temporal processing in F-344 rats. *Neuroscience*, *192*, 619–630.
- Peelle, J. E., & Wingfield, A. (2005). Dissociations in perceptual learning revealed by adult age differences in adaptation to time-compressed speech. *Journal of Experimental Psychology: Human Perception and Performance*, *31*, 1315.

- Pichora-Fuller, M., Schneider, B., & Daneman, M. (1995). How young and old adults listen to and remember speech in noise. *Journal of the Acoustical Society of America*, *97*, 593–608.
- Pichora-Fuller, M. K., & Levitt, H. (2012). Speech comprehension training and auditory and cognitive processing in older adults. *American Journal of Audiology*, *21*, 351.
- Puckett, A. C., Pandya, P. K., Moucha, R., Dai, W., & Kilgard, M. P. (2007). Plasticity in the rat posterior auditory field following nucleus basalis stimulation. *Journal of Neurophysiology*, *98*, 253–265.
- Phillips, S. L., Gordon-Salant, S., Fitzgibbons, P. J., & Yeni-Komshian, G. (2000). Frequency and temporal resolution in elderly listeners with good and poor word recognition. *Journal of Speech, Language, and Hearing Research*, *43*, 217–228.
- Rudner, M., Rönnerberg, J., & Lunner, T. (2011). Working memory supports listening in noise for persons with hearing impairment. *Journal of the American Academy of Audiology*, *22*, 156–167.
- Sabin, A., Clark, C., Eddins, D., & Wright, B. (2013). Different patterns of perceptual learning on spectral modulation detection between older hearing-impaired and younger normal-hearing adults. *Journal of the Association for Research in Otolaryngology*, *14*, 283–294.
- Salthouse, T. (1996). Constraints on theories of cognitive aging. *Psychonomic Bulletin & Review*, *3*, 287–299.
- Skoe, E., & Kraus, N. (2010). Auditory brain stem response to complex sounds: A tutorial. *Ear & Hearing*, *31*, 302–324.
- Smith, G. E., Housen, P., Yaffe, K., Ruff, R., Kennison, R. F., Mahncke, H. W., et al. (2009). A cognitive training program based on principles of brain plasticity: Results from the improvement in memory with plasticity-based adaptive cognitive training (IMPACT) study. *Journal of the American Geriatrics Society*, *57*, 594–603.
- Song, J. H., Skoe, E., Banai, K., & Kraus, N. (2012). Training to improve hearing speech in noise: Biological mechanisms. *Cerebral Cortex*, *22*, 1180–1190.
- Song, J. H., Skoe, E., Wong, P. C. M., & Kraus, N. (2008). Plasticity in the adult human auditory brainstem following short-term linguistic training. *Journal of Cognitive Neuroscience*, *20*, 1892–1902.
- Tremblay, K., Piskosz, M., & Souza, P. (2003). Effects of age and age-related hearing loss on the neural representation of speech cues. *Clinical Neurophysiology*, *114*, 1332–1343.
- Tremblay, K. L., Shahin, A. J., Picton, T., & Ross, B. (2009). Auditory training alters the physiological detection of stimulus-specific cues in humans. *Clinical Neurophysiology*, *120*, 128–135.
- VanderWerff, K. R., & Burns, K. S. (2011). Brain stem responses to speech in younger and older adults. *Ear & Hearing*, *32*, 168–180.
- Walton, J. P., Frisina, R. D., & O'Neill, W. E. (1998). Age-related alteration in processing of temporal sound features in the auditory midbrain of the CBA mouse. *The Journal of Neuroscience*, *18*, 2764–2776.
- Wang, H., Turner, J. G., Ling, L., Parrish, J. L., Hughes, L. F., & Caspary, D. M. (2009). Age-related changes in glycine receptor subunit composition and binding in dorsal cochlear nucleus. *Neuroscience*, *160*, 227–239.
- White-Schwoch, T., Carr, K. W., Anderson, S., Strait, D. L., & Kraus, N. (2013). Older adults benefit from music training early in life: Biological evidence for long-term training-driven plasticity. *The Journal of Neuroscience*, *33*, 17667–17674.
- Wilson, R. H., McArdle, R. A., & Smith, S. L. (2007). An evaluation of the BKB-SIN, HINT, QuickSIN, and WIN materials on listeners with normal hearing and listeners with hearing loss. *Journal of Speech, Language & Hearing Research*, *50*, 844–856.
- Wong, P. C. M., Ettlinger, M., Sheppard, J. P., Gunasekera, G. M., & Dhar, S. (2010). Neuroanatomical characteristics and speech perception in noise in older adults. *Ear & Hearing*, *31*, 471–479.
- Wong, P. C. M., Jin, J. X., Gunasekera, G. M., Abel, R., Lee, E. R., & Dhar, S. (2009). Aging and cortical mechanisms of speech perception in noise. *Neuropsychologia*, *47*, 693–703.
- Woodcock, R. W., McGrew, K. S., & Mather, N. (2001). *Woodcock-Johnson III tests of cognitive abilities*. Itasca, IL: Riverside Publishing.
- Zhu, J., & Garcia, E. (1999). *The Wechsler Abbreviated Scale of Intelligence (WASI)*. New York: Psychological Corporation.
- Zinke, W., Roberts, M., Guo, K., McDonald, J., Robertson, R., & Thiele, A. (2006). Cholinergic modulation of response properties and orientation tuning of neurons in primary visual cortex of anaesthetized Marmoset monkeys. *European Journal of Neuroscience*, *24*, 314–328.